Purpose of the Report

Enclosed is the 2011 Dialysis Facility Report (DFR) for your facility, based on data from the Centers for Medicare & Medicaid Services (CMS).

This DFR includes data specific to CCN(s): 492635

These data could be useful in quality improvement and assurance activities. The information contained in this report facilitates comparisons of patient characteristics, treatment patterns, transplantation rates, hospitalization rates, and mortality rates to local and national averages. Some of these comparisons account for the patient mix at this facility, including age, sex, race, and diabetic status. This report is provided as a resource for characterizing selected aspects of clinical experience at this facility relative to other caregivers in this state, ESRD Network, and across the United States.

In September 2011, each state's surveyor(s) will receive the DFRs with comments (if applicable) for all dialysis facilities in their state.

This report also provides you with advance notice of the updated quality measures (urea reduction ratio, hemoglobin, and patient survival) for your facility that will be reported on the Dialysis Facility Compare (DFC) website in November 2011 (www.medicare.gov).

Collaborators

CMS has contracted with the University of Michigan Kidney Epidemiology and Cost Center (UM-KECC) and Arbor Research Collaborative for Health to produce the *2011 Dialysis Facility Reports*.

How to Submit Comments

Between July 15, 2011 and August 15, 2011, you may submit comments to (1) CMS on the three DFC measures, (2) your state surveyor, or (3) UM-KECC. Please visit www.DialysisReports.org, log on to view your report, and click on the **Comments & Inquiries** tab. If you have questions after the comment period is over, please contact us directly at Support@DialysisReports.org or 1-877-665-1680.

- (1) **Dialysis Facility Compare:** Comment on the three DFC measures (see page 2) which will be reported on the DFC public website in November 2011. The comment period begins July 15, 2011 and ends August 15, 2011. Your comments will be shared with CMS but will **not** appear on the DFC website.
- (2) **State Surveyor:** Comment on your DFR for your state's surveyors. The surveyor(s) in your state will receive a copy of your DFR in September 2011 with your comments.
- (3) **UM-KECC:** Submit questions about your DFR to UM-KECC. You can also submit your suggestions to improve the DFR. These comments will not be shared with CMS or your state surveyor.

ADVANCED DIALYSIS CENTER - POTOMAC State: VA Network: 05 CCN: 492635

Dear Dialysis Facility Director:

This report has been prepared for this facility by the University of Michigan Kidney Epidemiology and Cost Center (UM-KECC) with funding from the Centers for Medicare & Medicaid Services (CMS). It is the sixteenth in a series of annual reports. This is one of 5,877 reports that have been sent to the ESRD Networks for distribution to ESRD providers in the U.S. Your state survey agency will receive this report in September 2011. Selected highlights from this report are given here. The information specific to this facility is printed in **bold type** for easy identification.

What's New This Year: As part of a continuing effort to improve the quality and relevance of this report for your facility, the following changes have been incorporated into your 2011 DFR. Emergency department and readmission statistics have been added to Table 2. In addition, most of the hospitalization summaries (Table 2) are now calculated for a four-year period. Values for each year from 2007-2010, along with a summarized statistic for the four-year period are reported for Medicare dialysis patients. We also reported the averages in your state, your ESRD Network, and the nation for this combined four-year period. We've added a new table that reports influenza vaccination summary statistics identified on Medicare claims for Medicare dialysis patients treated on December 31st of each year in your facility (Table 5). Average values for 2010 are also reported among patients in your state, Network, and the U.S. A new section on the prescription of erythropoiesis stimulating agents (ESA) and iron has been added to Table 6.

Dialysis Facility Compare Preview: Anemia management is reported as two measures: the percent of patients with hemoglobin values of less than 10 g/dL and the percent of patients with hemoglobin values greater than 12 g/dL. The URR and hemoglobin measures were calculated for Medicare approved dialysis facilities operating at any time during 2010. The hemoglobin measures were calculated only for patients treated with erythropoiesis stimulating agents (ESA). The patient survival measure was calculated for Medicare approved dialysis facilities operating at any time from 2007 through 2010. The following measures for this facility will appear on the DFC website:

	Measure Name	This Facility
1.	The percentage of Medicare hemodialysis patients treated in this facility during 2010 with URR >= 65% Number of patients included in calculation: 45	93%
2.	The percentage of Medicare patients treated in this facility during 2010 with ESA-treated hemoglobin <10 g/dL with ESA-treated hemoglobin >12 g/dL Number of patients included in calculation: 56	18% 9%
3.	Patient survival reported as "as expected," "better than expected," or "worse than expected" for the time period 2007-2010 for this facility Standardized Mortality Ratio (SMR): 2.53 P-value: <0.01	Worse than Expected

Please see Table 6 for more information on URR and ESA-treated hemoglobin for this facility. URR and ESA-treated hemoglobin measures based on 10 or fewer patients will be reported as "not available" on DFC. Table 1 provides additional information on patient survival. If the Standardized Mortality Ratio (SMR) is less than 1.00 and statistically significant (p<0.05), the patient survival classification is "Better than Expected" on DFC. If the facility SMR is greater than 1.00 and statistically significant (p<0.05), the patient survival classification is "Worse than Expected" on DFC. Otherwise, the patient survival classification is "As Expected" on DFC. Please note that the classification is not reported for a facility if the SMR is based on 3 or fewer expected deaths.

ADVANCED DIALYSIS CENTER - POTOMAC State: VA Network: 05 CCN: 492635

Overview: This report includes summaries of patient characteristics, treatment patterns, and patient outcomes for chronic dialysis patients who were treated in this facility between January 2007 and December 2010. Mortality, hospitalization, and transplantation statistics are reported for a three- or four-year period. Regional and national averages are included to allow for comparisons. Several of the summaries of patient mortality, hospitalization, and transplantation are adjusted to account for the characteristics of the patient mix at this facility, such as age, sex, race, ethnicity, and diabetes as a cause of ESRD. Unless otherwise specified, data refer to hemodialysis and peritoneal dialysis patients combined.

Summary data about the percent of patients with URR of 65% or higher (Table 6) and with central catheters (Table 7) are included, as suggested by Kidney Disease Outcomes Quality Initiative (KDOQI) Clinical Practice Guidelines. These practice pattern measures are strongly correlated with the mortality and hospitalization measures found in these reports. Note that elevated mortality or hospitalization rates may be due to a variety of causes unrelated to clinical practices, so it may not always be possible to identify clinical practices that explain those rates.

Mortality: Mortality summaries are provided in Table 1. Standardized Mortality Ratios (SMR) for all patients and for patients in their first year of ESRD treatment are calculated for each facility. Although mortality statistics can vary substantially from year to year, they are reported here as a key outcome for dialysis patients. The SMR compares the observed death rate in this facility to the death rate that would be expected based on national death rates for patients with the characteristics of the patients at this facility. For the overall SMR, time at risk and deaths within 60 days after transfer out of this facility are attributed to this facility. The first year SMR includes all incident patients at the facility for their entire first year of dialysis whether they transfer out of the facility or not. Time at risk and deaths after transplantation are excluded from the analysis. The SMR is adjusted for patient age, sex, race, ethnicity, diabetes as a cause of ESRD, duration of ESRD, nursing home status, body mass index (BMI) at incidence, comorbidities at incidence, and state population death rates. Additionally, each year's estimate is compared to the US mortality rates for the same year. The rate of withdrawal from dialysis and the percentage of deaths due to infection and cardiac related deaths are reported in the table to help in the interpretation of the mortality outcomes.

There was a 78% annual observed death rate among the patients treated at this facility during 2007-2010, while a rate of 31% would be expected after adjustment for the factors listed above. The SMR of observed to expected deaths is 2.53, which is 153% more deaths than expected at this facility. Among all U.S. facilities, 99% of facilities had an SMR lower than 2.53. This difference is statistically significant (p<0.05), so this higher mortality is unlikely to be due to random chance and probably represents a real difference from the expected mortality in the nation.

There was a 71% annual observed first year death rate among the patients starting dialysis at this facility during 2007-2009, while a rate of 27% would be expected after adjustment for the factors listed above. The first year SMR of observed to expected deaths is 2.61, which is 161% more deaths than expected at this facility. Among all U.S. facilities, 98% of facilities had an SMR lower than 2.61. This difference is statistically significant (p<0.05), so this higher mortality is unlikely to be due to random chance and probably represents a real difference from the expected mortality in the nation.

Hospitalization: Hospitalization summaries are reported for Medicare patients in Table 2. The table includes information on the number of days hospitalized, the number of hospital admissions and the diagnoses present at admission, and the number of emergency department (ED) visits for patients at this facility. The Standardized Hospitalization Ratio (SHR) for Days, Admissions, and ED visits compares the observed number of events (be it days hospitalized, admissions, or ED visits) to the number that would be expected based on national rates for patients with the characteristics of the patients at this facility. The three measures are adjusted for age, sex, diabetes, duration of ESRD, nursing home status, BMI, and comorbidities at incidence. Additionally, each year's estimate is compared to the US rates for the same year. As in the mortality calculation, time at risk and events within 60 days after transfer out of this facility are attributed to this facility. Time at risk and events starting 3 days before transplantation are excluded from the analysis.

The SHR (Days) of observed to expected number of days hospitalized at this facility during 2007-2010 is 1.54, which is 54% higher than expected. The SHR (Admissions) of observed to expected number of admissions for patients at this facility during 2007-2010 is 1.17, which is close to the national reference value. The SHR (ED) of observed to expected number of ED visits for patients at this facility during 2007-2009 is 0.95, which is close to the national reference value.

Infection: Information on hospitalizations for septicemia reported in Table 2 is based on Medicare patient claims.

ADVANCED DIALYSIS CENTER - POTOMAC State: VA Network: 05 CCN: 492635

The percentage of Medicare dialysis patients at this facility hospitalized with septicemia during 2007-2010 was 24%, compared to 11% nationally. It may be useful to review infection control at this facility.

Transplantation: Transplantation summaries are reported in Table 3. The Standardized Transplantation Ratio (STR) represents relative first transplantation rates (observed/expected) for patients at this facility adjusting for patient age. Additionally, each year's estimate is compared to the US transplantation rates for the same year. Patients who are 70 or older as well as those with a prior kidney transplant are excluded. As in the mortality and hospitalization calculations, time at risk and transplants within 60 days after transfer out of this facility are attributed to this facility.

Of the patients under age 70 treated at this facility during 2007-2010 who had not previously received a transplant, none were transplanted annually, while a rate of 4% would be expected for these patients. The STR of observed to expected number of patients transplanted for this facility is 0.00, which is 100% lower than expected for this facility. This difference is not statistically significant (p>=0.05) and is plausibly due to random chance.

Transplant Waitlist: Table 4 summarizes waitlist information for patients under age 70 being treated at this facility at the end of each year. Unlike the mortality, hospitalization, and transplantation statistics, the waitlist statistics do not include patients who transferred out of the facility.

Among the 42 dialysis patients under age 70 treated at this facility on December 31, 2010, 19% were on the kidney transplant waitlist compared to 24% nationally. This difference is not statistically significant (p>=0.05) and is plausibly due to random chance.

Influenza Vaccination: Table 5 summarizes influenza vaccination information for Medicare patients being treated at this facility at the end of each year. Similar to the waitlist table, influenza vaccination statistics do not include patients who transferred out of the facility.

Among the 47 Medicare dialysis patients treated at this facility on December 31, 2009, 49% were vaccinated between September 1 and December 31, 2009 compared to 69% nationally. This difference is statistically significant (p<0.05) and is unlikely to be due to random chance.

Practice Patterns: Table 6 summarizes the dialytic modality, hemoglobin, and URR for patients treated at this facility during each year. These data are derived from CMS Medicare paid dialysis claim data. Vascular access data for prevalent and incident patients as reported by the CMS Fistula First project are summarized in Table 7.

There were 56 ESA-treated dialysis patients included in the analysis of Medicare claims data of hemoglobin for 2010 at this facility. The average hemoglobin calculated for these patients is 10.9 g/dL, compared to 11.4 g/dL nationally. Among the 45 hemodialysis patients in this facility included in the analysis of Medicare claims data of URR in 2010, 93% had URR above the KDOQI minimum value for URR (URR \geq 65%), compared to 97% nationally.

We do not have Fistula First information from this facility in 2010. See Tables 6 and 7 for more information about practice patterns.

Patient Characteristics: Characteristics of patients starting dialysis during 2007-2010 are reported in Table 8. Table 9 gives summaries for all dialysis patients being treated at the end of each year, 2007-2010. Comorbidities are reported in Table 10 for Medicare dialysis patients being treated at the end of each year, 2007-2009.

There were 46 patients with Medical Evidence Forms (CMS-2728) which indicated that they started treatment at this facility during 2010. The average number of comorbidities reported for new patients is 3.0, which is lower than the average of 3.2 reported nationally. Also, 33% of these patients were not under the care of a nephrologist before starting dialysis, compared to 30% nationally. Furthermore, 100% of these patients were informed of their transplant options, compared to 77% nationally. The average serum albumin calculated for these patients (before first dialysis) is 2.7 g/dL, which is lower than the national average value of 3.1 g/dL. The average residual renal function (GFR), estimated by the 4-variable MDRD equation, was 13.9 mL/min, which is higher than the national average value of 11.2 mL/min.

Among patients treated at this facility on December 31, 2010, 75% were treated in a nursing home during the year, which is higher than the national average value of 13%. The average number of comorbidities reported on Medicare claims in 2009 for Medicare patients in this facility on December 31, 2009 is 7.2, which is higher than the national average value of 4.2.

ADVANCED DIALYSIS CENTER - POTOMAC State: VA Network: 05 CCN: 492635

Facility Information: General information about this facility is provided in Tables 12, 13, and 14. Table 12 provides counts of patients treated, Medicare eligibility, treatment modality, and staffing from the Annual Facility Survey (Form CMS-2744). Table 13 reports survey and certification activity. Table 14 includes services provided by this facility as well as information on ownership.

This facility reported having 16 staff members to care for 71 patients at the end of 2010. This facility reported having 9 stations available as of July 31, 2011. Additional information regarding patient counts, patient modality, and facility staffing is available in Table 12.

Sources of Patient Data: This report is based primarily on Medicare claims and data collected for CMS. Patients were assigned to this facility based on the Standard Information Management System (SIMS) database, Medicare claims, and Medical Evidence Forms (Form CMS-2728). Network 05 has a list of the patients included in the mortality analyses for this facility. Table 10 reports comorbidities reported on Medicare claims. Table 11 reports the number of patients placed in this facility for analyses in Tables 1, 2, 3, 9, and 10 of this report. Table 11 also provides information on patient status at the end of the year, both for patients who remain in this facility until the end of each year, and for those who transfer out of this facility, receive a transplant, or die during the year.

These are just a few highlights of the statistics you will find in this report based on the data for this facility. We hope that this report is of interest to you and that you will discuss it with your staff. We welcome any questions or comments you might have about the content of the current report or any suggestions you might have for future reports. Comments or suggestions can be submitted via the secure site www.DialysisReports.org between July 15, 2011 and August 15, 2011. If you have questions after the comment period is over, please contact us directly at Support@DialysisReports.org or 1-877-665-1680.

For a complete description of the methods used to calculate the statistics in this report, please see the *Guide to the 2011 Dialysis Facility Reports*. The *Guide* is available from ESRD Network 05, and is also on the Dialysis Reports web site at www.DialysisReports.org.

Prepared by

The University of Michigan Kidney Epidemiology and Cost Center (UM-KECC) under contract with the Centers for Medicare & Medicaid Services

TABLE 1: Mortality Summary for All Dialysis Patients (2007-10) & New Dialysis Patients (2007-09)*1

			,	This Faci	lity			onal Average Year, 2007-20	
	Measure Name	2007	2008	2009	2010	2007-2010	State	Network	U.S.
All	Patients: Death Rates								
1a	Patients (n=number)	73	93	100	133	399 *8	83.8	86.2	92.4
1b	Patient years (PY) at risk (n)	26.2	33.2	39.7	51.8	150.8 *8	58.8	60.1	61.4
1c	Deaths (n)	18	30	35	35	118 *8	11.8	12.2	12.1
1d	Expected deaths (n)	9.2	11.3	12.6	13.4	46.6 *8	10.7	11.1	12.1
1e	Death rate per 100 PY (% of 1b)	68.8	90.3	88.2	67.6	78.2	20.1	20.3	19.7
1f	Expected death rate per 100 PY (% of 1b)	35.3	34.2	31.8	25.9	30.9	18.2	18.4	19.7
All	Patients: Categories of Death								
1g	Withdrawal from dialysis prior to death (% of 1c)	5.6	20.0	17.1	28.6	19.5	25.5	20.7	25.0
1h	Death due to: Infections (% of 1c)	0.0	6.7	5.7	2.9	4.2	17.6	15.6	16.1
	Cardiac causes (% of 1c)	5.6	0.0	2.9	0.0	1.7	27.5	24.1	25.6
1i	Dialysis unrelated deaths *3 (n; excluded from SMR)	0	0	0	0	0 *8	0.1	0.1	0.1
All	Patients: Standardized Mortality Ratio (SMR)								
1j	SMR *4	1.95	2.64	2.78	2.61	2.53	1.10	1.10	1.00
1k	P-value*5	0.013	< 0.01	< 0.01	< 0.01	< 0.01	n/a	n/a	n/a
11	Confidence interval for SMR *6								
	High (95% limit)	3.08	3.77	3.86	3.64	3.03	n/a	n/a	n/a
	Low (5% limit)	1.16	1.78	1.93	1.82	2.10	n/a	n/a	n/a
1m	SMR Percentiles for this Facility (i.e. percent of facilities	es with lower:	mortality r	rates)*7					
	In this State	95	99	98	98	96			
	In this Network	93	99	98	98	97			
	In the U.S.	96	99	99	98	99			
NT.	- Dedicator Electory - Deadly Dedica	2007	2000	2000		2000	Reg	gional Avera	1ges
	w Patients: First Year Death Rates	2007 29	2008	2009	2	2007-2009		Year, 2007-2	
1n	New Patients (n=number)		42	40		111 *8	18.3	18.9	19.1
10	Patient years (PY) at risk (n)	21.0	29.4	26.7		77.1 *8	15.7	16.1	16.4
1p	Deaths (n)	13	21	21		55 *8	4.1	4.4	4.4
lq	Expected deaths (n)	5.7	8.6	6.8		21.0 *8	3.8	4.0	4.4
lr	Death rate per 100 PY	61.9 27.2	71.4 29.1	78.8 25.3		71.3	25.9 24.4	27.2 24.6	26.7 26.7
1s	Expected death rate per 100 PY	21.2	29.1	23.3		27.3	24.4	24.0	20.7
	w Patients: Categories of Deaths		10.0	0.7		10.5	22.0	20.7	
1t	Withdrawal from dialysis prior to death (% of 1r)	7.7	19.0	9.5		12.7	33.0	29.5	34.2
lu	Death due to: Infections (% of 1r)	7.7	9.5	9.5		9.1	23.8	20.9	22.9
	Cardiac causes (% of 1r)	0.0	9.5	0.0		3.6	30.3	30.3	32.9
Ne	w Patients: First Year Standardized Mortality R	atio (SMR)							
1v	SMR *4	2.27	2.45	3.11		2.61	n/a	n/a	n/a
1w	P-value*5	0.012	< 0.01	< 0.01		< 0.01	n/a	n/a	n/a
1x	Confidence interval for SMR*6								
	High (95% limit)	3.89	3.75	4.75		3.40	n/a	n/a	n/a
	Low (5% limit)	1.21	1.52	1.93		1.97	n/a	n/a	n/a
1y	First Year SMR Percentiles for this Facility (i.e. percent	of facilities w	ith lower r		ates)*7				
	In this State	94	97	97		97			
	In this Network	91	94	97		96			
	In the U.S.	93	94	97		98			

n/a = not applicable

[*1] See *Guide, Section IV.*[*2] Values are shown for the average facility, annualized.

[*3] Defined as deaths due to street drugs and accidents unrelated to treatment.

[*4] Calculated as a ratio of deaths to expected deaths(1c to 1d for all patients, 1p to 1q for new patients); not shown if there are fewer than 3 expected deaths.

[*5] A p-value less than 0.05 indicates that the difference between the actual and expected mortality is probably real and is not due to random chance alone, while a p-value greater than or equal to 0.05 indicates that the difference could plausibly be due to random chance.

^[*6] The confidence interval range represents uncertainty in the value of the SMR due to random variation.
[*7] All facilities are included in ranking, regardless of the number of expected deaths.
[*8] Sum of 4 years (all patients), or 3 years (new patients), used for calculations; should not be compared to regional averages.

TABLE 2: Hospitalization Summary for Medicare Dialysis Patients *1 , 2007-2010

			Т	his Facili			onal Average Year, 2007-20		
	Measure Name	2007	2008	2009	2010	2007-2010	State	Network	U.S.
Me	edicare Dialysis Patients								
2a	Medicare dialysis patients (n)	65	82	86	113	346 *6	69.9	70.2	72.2
2b	Patient years (PY) at risk (n)	21.0	28.1	33.0	42.0	124.2 *6	47.6	47.1	44.8
Da	ys Hospitalized Statistics								
2c	Total days hospitalized (n)	938	1046	1188	934	4106 *6	603.6	665.9	650.7
2d	Expected total days hospitalized (n)	487.6	635.0	751.1	801.2	2674.8 *6	672.1	669.0	651.0
2e	Days hospitalized per PY	44.6	37.2	36.0	22.2	33.1	12.7	14.1	14.5
2f	Expected days hospitalized per PY	23.2	22.6	22.8	19.1	21.5	14.1	14.2	14.5
2g	Standardized Hospitalization Ratio (Days)*3	1.92	1.65	1.58	1.17	1.54	0.90	1.00	1.00
2h	Percentiles for this Facility (i.e. % of facilities with lower hospitalize	ation rates	[days])*4						
	In this State	98	96	95	86	97	n/a	n/a	n/a
	In this Network	95	94	91	71	92	n/a	n/a	n/a
	In the U.S.	96	93	92	71	92	n/a	n/a	n/a
Λd	mission Statistics			, -	, -	7-			
Au 2i	Total admissions (n)	68	91	113	107	379 *6	83.9	95.5	88.3
2j	Expected total admissions (n)	56.1	74.3	91.2	102.6	324.2 *6	91.6	91.0	88.4
2 _J 2k	Admissions per PY	3.2	3.2	3.4	2.5	3.1	1.8	2.0	2.0
21	Expected admissions per PY	2.7	2.6	2.8	2.4	2.6	1.9	1.9	2.0
2m	Standardized Hospitalization Ratio (Admissions) *3	1.21	1.23	1.24	1.04	1.17	0.92	1.05	1.00
	•				1.04	1.17	0.72	1.03	1.00
2n	Percentiles for this Facility (i.e. % of facilities with lower hospitalized by the facilities of the facilities with lower hospitalized by the facilities wi				7.0	97	,	,	,
	In this State	80	92	91	76	87	n/a	n/a	n/a
	In this Network	68	75	76	56	66	n/a	n/a	n/a
	In the U.S.	78	78	80	59	74	n/a	n/a	n/a
20	Diagnoses Associated with Hospitalization (% of 2a)*5								
	Septicemia	24.6	26.8	31.4	16.8	24.3	11.8	13.5	11.2
	Acute myocardial infarction	3.1	7.3	1.2	0.9	2.9	4.5	4.8	4.1
	Congestive heart failure	20.0	15.9	22.1	22.1	20.2	23.7	26.8	21.5
	Cardiac dysrhythmia	15.4	12.2	22.1	9.7	14.5	12.8	13.9	12.0
	Cardiac arrest	0.0	2.4	1.2	1.8	1.4	1.4	1.3	1.3
2p	One day admissions (% of 2i)	4.4	5.5	8.8	6.5	6.6	12.2	13.9	13.4
2q	Average length of stay (days per admission; 2c/2i)	13.8	11.5	10.5	8.7	10.8	7.2	7.0	7.4
2r	Readmissions within 30 days (n)	26	21	36	34	117 *6	99.0	126	110
2s	Admissions that result in readmission within 30 days (% of 2i)	38.2	23.1	31.9	31.8	30.9	29.5	33.1	31.1
							Res	gional Avera	ages
En	nergency Department(ED) Statistics	2007	2008	2009	20	007-2009	Per	Year, 2007-	2009*2
2t	Total ED visits (n)	75	95	126		296 *6	438.3	441.2	385.7
2u	Expected total ED visits (n)	77.6	101	131		310 *6	404.2	400.2	385.8
2v	ED visits per PY	3.6	3.4	3.8		3.6 *6	3.1	3.1	2.8
2w	Expected ED visits per PY	3.7	3.6	4.0		3.8 *6	2.8	2.8	2.8
2x	Standardized Hospitalization Ratio (ED)*3	0.97	0.94	0.96		0.95	1.08	1.10	1.00
2y	Percentiles for this Facility (i.e. % of facilities with lower hospitalize	ation rates	[ED])*4						
	In this State	29	31	36		32	n/a	n/a	n/a
	In this Network	32	30	34		30	n/a	n/a	n/a
	In the U.S.	49	44	47		45	n/a	n/a	n/a
2z	Patients with ED visit (% of 2a)	61.5	59.8	62.8		61.4*7	64.5	64.6	57.9
2aa	ED visits that result in hospitalization (% of 2t)	76.0	77.9	73.8		75.7	42.9	51.7	50.6
2ab	Admissions that originate in the ED (% of 2i)	83.8	81.3	82.3		82.4*7	74.1	78.2	71.6

 $n/a = not \ applicable.$

n/a = not applicable.

[*1] Based on patients with Medicare as primary insurer; see *Guide, Section V*.

[*2] Values are shown for the average facility, annualized.

[*3] Calculated as ratio of actual to expected events (2e/2f for days, 2k/2l for admissions, and 2v/2w for ED visits); not shown if there are fewer than 5 patient years at risk.

[*4] All facilities are included in ranking, regardless of the number of patient years at risk.

[*5] Includes diagnoses present at admissions and diagnoses added during the hospital stay.

[*6] Sum of 4 years (days and admissions) or 3 years (ED visits), used for calculations; should not be compared to regional averages.

[*7] Percent is based on the sum of three years.

TABLE 3: Transplantation Summary for Dialysis Patients under Age 70*1, 2007-2010

				This Faci	lity		Regional Averages *2, per Year, 2007-2010			
	Transplants (n) Donor type (sums to 3b *3) Living Donor (n) Deceased Donor (n) ents who have not Previously Received a Tr Eligible patients (n) Patient years (PY) at risk (n) First transplants *4 (n) Expected First transplants (n) First transplant rate per 100 PY Expected First transplant rate per 100 PY Donor type (sums to 3f *5) Living Donor (n) Deceased Donor (n) adardized 1st Transplantation Ratio (STR) * STR *7 P-value *8 95% Confidence interval for STR *9 Upper limit Lower limit STR Percentiles for this Facility (i.e. percent of facility (i.e. percent of facility (i.e. percent)	2007	2008	2009	2010	2007-2010	State	Network	U.S.	
3a	Eligible patients (n)	41	47	58	75	221 *11	58.9	60.2	63.4	
3b	Transplants (n)	0	0	0	0	0 *11	2.5	2.4	2.3	
3c	Donor type (sums to 3b*3)									
	Living Donor (n)	0	0	0	0	0 *11	0.8	0.8	0.7	
	Deceased Donor (n)	0	0	0	0	0 *11	1.7	1.7	1.6	
Pa	tients who have not Previously Received a Tr	ansplant								
3d	Eligible patients (n)	39	46	56	71	212 *11	53.5	54.7	57.0	
3e	Patient years (PY) at risk (n)	14.7	16.4	23.7	33.0	87.7 *11	37.9	38.6	38.4	
3f	First transplants *4 (n)	0	0	0	0	0 *11	2.2	2.1	2.0	
3g	Expected First transplants (n)	0.6	0.6	1.0	1.3	3.5 *11	1.9	2.0	2.0	
3h	First transplant rate per 100 PY	0.0	0.0	0.0	0.0	0.0	5.8	5.5	5.1	
3i	Expected First transplant rate per 100 PY	3.9	3.8	4.1	4.0	4.0	5.0	5.1	5.1	
3j	Donor type (sums to 3f*5)									
	Living Donor (n)	0	0	0	0	0 *11	0.6	0.6	0.6	
	Deceased Donor (n)	0	0	0	0	0 *11	1.5	1.5	1.4	
Sta	andardized 1st Transplantation Ratio (STR)	*6								
3k	STR *7					0.00	1.15	1.08	1.00	
31	P-value*8					0.061	n/a	n/a	n/a	
3m	95% Confidence interval for STR *9									
	Upper limit	•			•	1.06	n/a	n/a	n/a	
	Lower limit				ė	0.00	n/a	n/a	n/a	
3n	STR Percentiles for this Facility (i.e. percent of faci	lities with lower tr	ansplantat	ion rates)*	10					
	In this State					7	n/a	n/a	n/a	
	In this Network		-			12	n/a	n/a	n/a	
	In the U.S.					10	n/a	n/a	n/a	

n/a = not applicable.

[*1] See *Guide, Section VI*.

[*2] Values are shown for the average facility, annualized.

[*3] Values may not sum to 3b due to unknown donor type.

[*4] Among first transplants that occurred after the start of dialysis from 2007-2010, 3.9% of transplants in the U.S. were not included because the transplant occurred less than 90 days after the start of ESRD and 0.9% were not included because the patient was not assigned to a facility at time of transplant.

[*5] Values may not sum to 3f due to unknown donor type.

[*6] This section is not reported if there are fewer than 3 expected transplants.

[*7] Standardized Transplantation Ratio calculated as ratio of actual (30) to expected (3e) transplants.

^[*8] Standardized Transplantation Ratio calculated as ratio of actual (3f) to expected (3g) transplants.

[*8] A p-value less than 0.05 indicates that the difference between the actual and expected transplants is probably real and is not due to random chance, while a p-value greater than or equal to 0.05 indicates that the difference is plausibly due to random chance.

^[*9] The confidence interval range represents uncertainty in the value of the STR due to random variation. [*10] All facilities are included in ranking, regardless of the number of expected transplants. [*11] Sum of 4 years used for calculations; should not be compared toregional averages.

TABLE 4: Waitlist Summary for Dialysis Patients under Age 70 Treated as of December 31st of Each Year*1, 2007-2010

			This Fa	cility		Regional Averages *2, 2010			
	Measure Name	2007	2008	2009	2010	State	Network	U.S.	
a	Eligible patients on 12/31 (n)	21	27	38	42	48.5	48.1	46.	
b	Patients on the waitlist (% of 4a)	0.0	22.2	28.9	19.0	26.8	22.9	23.	
c	P-value (compared to U.S. value) *3	< 0.01	0.519	0.304	0.304	n/a	n/a	n	
d	Patients on the waitlist by subgroup (% of corresponding value in 4e)								
	Age < 40		0.0	0.0	50.0	39.3	32.5	35	
	Age 40-69	0.0	23.1	30.6	17.5	25.0	21.5	22	
	Male	0.0	27.8	28.6	13.3	26.3	22.2	23	
	Female	0.0	11.1	29.4	33.3	27.4	23.7	23	
	African American	0.0	23.1	29.2	17.9	27.2	22.6	21	
	Asian/Pacific Islander	0.0				43.8	34.9	35	
	Native American		50.0	100	100	37.5	33.3	17	
	White, Hispanic	•		0.0	0.0	25.8	24.6	26	
	White, Non-Hispanic	0.0	18.2	18.2	16.7	24.1	22.2	23	
	Other/unknown race	0.0	0.0	100		31.3	20.9	25	
	Diabetes	0.0	23.1	28.6	31.6	22.4	19.1	19	
	Non-diabetes	0.0	21.4	29.2	8.7	29.7	25.2	27	
	Previous kidney transplant	0.0		0.0	0.0	45.2	43.3	46	
	No previous kidney transplant	0.0	22.2	29.7	21.1	24.9	20.6	21	
	< 2 years since start of ESRD	0.0	0.0	14.3	15.4	19.6	16.0	15	
	2-4 years since start of ESRD	0.0	25.0	40.0	23.1	30.6	24.9	27	
	5+ years since start of ESRD	0.0	50.0	35.7	18.8	31.4	28.8	29	
e	Eligible patients in 4a by subgroup (n)								
	Age < 40	0	1	2	2	5.9	5.9	6	
	Age 40-69	21	26	36	40	42.5	42.2	40	
	Male	12	18	21	30	27.4	27.3	26	
	Female	9	9	17	12	21.1	20.8	20	
	African American	9	13	24	28	30.5	31.3	19	
	Asian/Pacific Islander	2	0	0	0	1.5	1.2	2	
	Native American	0	2	1	1	0.1	0.1	(
	White, Hispanic	0	0	1	1	1.5	1.2	7	
	White, Non-Hispanic	9	11	11	12	14.4	13.8	16	
	Other/unknown race	1	1	1	0	0.4	0.4	C	
	Diabetes	11	13	14	19	19.4	18.5	20	
	Non-diabetes	10	14	24	23	29.1	29.6	25	
	Previous kidney transplant	1	0	1	4	4.5	4.7	4	
	No previous kidney transplant	20	27	37	38	43.9	43.3	42	
	< 2 years since start of ESRD	11	11	14	13	18.1	17.8	17	
	< 2 years since start of ESRD 2-4 years since start of ESRD	8	8	10	13	15.0	17.8	17	
	2 + years since start of Lond	o	О	10	13	15.0	14.0	14	

n/a = not applicable

[*1] See *Guide, Section VII.*[*2] Values are shown for the average facility.

[*3] Facility waitlist percentage is compared to the U.S. waitlist percentage for that year: 23.8% (2007), 24.0% (2008), 24.2% (2009), 23.8% (2010).

A p-value greater than 0.05 indicates that the difference between percent of patients waitlisted at the facility and national percentage is plausibly due to random chance.

TABLE 5: Influenza Vaccination Summary for Medicare Dialysis Patients Treated on December 31st of Each Year*1, Flu Seasons September 2007-March 2010

		Th	is Facility	•		onal Averag 009-2010*2	es,
	Measure Name	2007-08	2008-09 2	2009-10	State	Network	U.S.
5a	Eligible patients on 12/31 (n)	32	42	47	51.9	51.5	49.1
5b	Patients vaccinated between Sep. 1 and Dec. 31 (% of 5a)	15.6	33.3	48.9	76.5	71.4	69.0
5c	P-value *3 (for 5b compared to U.S. value *4)	< 0.01	< 0.01	< 0.01	n/a	n/a	n/a
5d	Patients vaccinated between Sep. 1 and Mar. 31 (% of 5a)	15.6	33.3	48.9	77.6	72.5	70.3
5e	P-value *3 (for 5d compared to U.S. value *5)	< 0.01	< 0.01	< 0.01	n/a	n/a	n/a
5f	Patients vaccinated between Sep. 1 and Dec. 31						
	by subgroup (% of corresponding value in 5g)						
	Age < 18			·	45.0	43.9	49.5
	Age 18-39		100	0.0	67.8	64.9	64.8
	Age 40-64	10.0	28.6	42.1	75.6	70.6	68.6
	Age 65-74	18.2	30.0	53.8	77.6	72.7	69.8
	Age 75+	18.2	35.3	57.1	80.7	73.9	70.8
	Male	14.3	26.3	52.4	76.7	71.8	69.3
	Female	16.7	39.1	46.2	76.4	71.0	68.7
	African American	11.8	40.9	51.6	74.8	70.0	65.4
	Asian/Pacific Islander	0.0	0.0	50.0	83.5	77.3	70.8
	Native American		50.0	0.0	83.3	74.1	73.2
	White	30.0	23.5	50.0	78.8	73.6	71.3
	Other/unknown race	0.0		0.0	75.0	66.0	65.0
	Hispanic	0.0	0.0	0.0	80.4	74.7	71.5
	< 1 year since start of ESRD	20.0	61.5	44.4	69.3	62.7	58.4
	1-2 years since start of ESRD	11.1	16.7	64.3	75.7	71.4	69.2
	3+ years since start of ESRD	15.4	23.5	41.7	79.2	74.1	72.4
5g	Eligible patients in 5a by subgroup (n)						
	Age < 18	0	0	0	< 1.0	< 1.0	< 1.0
	Age 18-39	0	1	1	4.0	3.6	3.9
	Age 40-64	10	14	19	23.0	22.6	21.5
	Age 65-74	11	10	13	13.1	13.3	12.1
	Age 75+	11	17	14	11.6	11.8	11.4
	Male	14	19	21	27.6	27.7	26.6
	Female	18	23	26	24.3	23.8	22.5
	African American	17	22	31	30.4	31.3	18.4
	Asian/Pacific Islander	4	1	2	1.7	1.3	2.1
	Native American	0	2	1	0.1	0.1	0.8
	White	10	17	12	19.4	18.5	27.3
	Other/unknown race	1	0	1	0.4	0.3	0.5
	Hispanic	1	1	2	1.3	1.0	7.1
	< 1 year since start of ESRD	10	13	9	8.4	8.5	8.6
	1-2 years since start of ESRD	9	12	14	15.0	14.5	14.3
	3+ years since start of ESRD	13	17	24	28.4	28.5	26.2

n/a = not applicable

[*1] Based on patients with Medicare as primary insurer; see *Guide, Section VIII*.

[*2] Values are shown for the average facility.

[*3] A p-value greater than or equal to 0.05 indicates that the difference between percent of patients vaccinated at the facility and national percentage is plausibly due to random chance.

[*4] Compared to the U.S. value for that year and time period (9/1-12/31): 63.3% (2007-08), 64.2% (2008-09), 69.0% (2009-10).

[*5] Compared to the U.S. value for that year and time period (9/1-3/31): 63.8% (2007-08), 64.9% (2008-09), 70.3% (2009-10).

TABLE 6: Facility Modality, Hemoglobin, and Urea Reduction Ratio for Medicare Dialysis Patients *1 , 2007-2010

			This Fa	cility		Regiona	l Averages *	² , 2010
	Measure Name	2007	2008	2009	2010	State	Network	U.S.
Mo	dality (among all dialysis patients with ESRD for	90+ days and 1+ cla	im at this	s facility)				
6a	Patients treated during year (n)	103	135	148	165	83.2	82.6	75.0
6b	Patients months treated during year (n) *3	373	460	541	639	621.7	604.6	558.3
6c	Modality (% of 6b; sums to 100%)							
	Hemodialysis	100	100	100	100	93.2	93.4	92.5
	CAPD/CCPD	0.0	0.0	0.0	0.0	5.8	5.8	6.6
	Other dialysis *4	0.0	0.0	0.0	0.0	1.0	0.9	0.9
6d	Percent of patient months prescribed ESA by Modality *5							
	Hemodialysis	100	100	100	100	100	100	100
	CAPD/CCPD					80.5	79.5	81.1
6e	Percent of patient months prescribed Iron by Modality *5							
	Hemodialysis	46.4	49.6	50.5	44.9	62.4	58.3	61.9
	CAPD/CCPD					18.2	15.0	15.1
He	moglobin (among ESA-treated dialysis patients wi	th ESRD for 90+ da	ys and 4	+ Hemog	lobin clair	ns at this fa	cility)	
6f	Eligible patients (n)	27	36	41	56	54.5	53.4	49.0
6g	Average hemoglobin (g/dL)	10.9	10.9	11.0	10.9	11.3	11.4	11.4
6h	Hemoglobin categories (% of 6f; sums to 100%)							
	< 10 g/dL	18.5	8.3	9.8	17.9	3.1	3.6	3.1
	10-11 g/dL	25.9	36.1	31.7	26.8	18.6	17.5	17.9
	11-12 g/dL	55.6	50.0	51.2	46.4	69.5	68.6	68.3
	> 12 g/dL	0.0	5.6	7.3	8.9	8.8	10.3	10.7
6i	Eligible hemodialysis (HD) patients (n)*6	27	36	41	56	52.0	50.9	46.5
6j	Hemoglobin categories among HD pts (% of 6i; sums to	100%)						
-	< 10 g/dL	18.5	8.3	9.8	17.9	3.3	3.7	3.2
	10-11 g/dL	25.9	36.1	31.7	26.8	17.9	16.9	17.4
	11-12 g/dL	55.6	50.0	51.2	46.4	69.9	69.0	68.6
	> 12 g/dL	0.0	5.6	7.3	8.9	8.9	10.5	10.8
6k	Eligible peritoneal dialysis (PD) patients (n)*6	0	0	0	0	3.3	3.2	3.1
61	Hemoglobin categories among PD pts (% of 6k; sums to							
	< 10 g/dL					8.4	8.6	8.3
	10-11 g/dL					29.1	27.9	27.1
	11-12 g/dL					54.2	53.1	51.7
	> 12 g/dL					8.3	10.4	12.9
Ur	ea Reduction Ratio (URR; among HD patients wit	h ESRD for 183+ d:	avs and 4	+ URR cl	laims at th	4	7	
6m	Eligible patients (n)	23	27	29	45	49.8	48.8	44.6
6n	URR categories (% of 6m; sums to 100%)					.,,,	.0.0	
011	< 60.0 %	0.0	0.0	0.0	0.0	1.1	1.1	1.1
	60.0-64.9 %	4.3	3.7	3.4	6.7	2.6	2.7	2.3
	65.0-69.9 %	8.7	14.8	3.4	8.9	12.5	13.2	10.8
	70.0-74.9 %	34.8	37.0	27.6	28.9	38.8	39.5	35.6
	75+ %	52.2	44.4	65.5	55.6	45.0	43.5	50.2
60								
60	URR 65+ (% of 6m; meets a KDOQI guideline) URR percentiles for this facility (i.e. % of facilities with a	95.7	96.3	96.6	93.3	96.2	96.2	96.6
бр	In this State	smaner percentage of p	yatients wi	ın UKK 65 42	+) 18	n/a	n/a	n/a
	In this Network	40	44	44	20	n/a n/a	n/a n/a	n/a n/a
	In the U.S.	35	38	35	16	n/a	n/a	n/a

n/a = not applicable

[*1] See *Guide, Section IX.*[*2] Values are shown for the average facility.

[*3] Patients may be counted up to 12 times per year.

[*4] Other dialysis includes patients who switch between HD and PD during the month and patients for whom modality is unknown or missing.

[*5] Percent of patient months represented by the corresponding modality percent in 6c.

[*6] Sum of eligible HD and PD patients may not add to 6f.

[*7] Claims identified as having 4 or more dialysis sessions per week were excluded from the URR calculations. Among eligible claims in the US, 1.33% were excluded due to frequent dialysis in 2007-2010.

TABLE 7: Vascular Access Information (CMS Fistula First)*1, 2007-2010

			This Fa	cility	Regiona	l Averages *2	, 2010	
	Measure Name	2007	2008	2009	2010	State	Network	U.S.
Va	scular Access							
7a	Prevalent hemodialysis patient months *3	0	0	0	0	n/a	n/a	n/a
7b	Vascular access type in use (% of 7a; sums to 100%)							
	Arteriovenous fistula					53.9	52.7	56.0
	Arteriovenous graft					21.4	23.1	20.4
	Catheter	·				24.6	24.1	23.5
	Other/Missing					0.0	0.0	0.1
7c	Arteriovenous fistulae in place (% of 7a)*4					61.7	60.5	63.8
7d	Catheter only \geq 90 days (% of 7a)*5					10.3	10.0	9.2
Va	scular Access at First Treatment							
7e	Incident hemodialysis patients (n)	0	0	0	0	16.5	16.3	17.0
7f	Vascular access type in use (% of 7e; sums to 100%)*6							
	Arteriovenous fistula					15.4	15.2	17.4
	Arteriovenous graft					7.0	9.3	7.7
	Catheter					77.6	75.4	74.7
	Other/Missing					0.0	0.1	0.2
7g	Arteriovenous fistulae in place (% of 7e)*4				•	31.6	32.3	34.9

n/a = not applicable

[*1] See *Guide, Section X*.

[*2] Values are shown for the average facility.

[*3] Patients may be counted up to 12 times per year.

[*4] Includes all patients with fistulae, regardless of whether or not they received their hemodialysis treatments using their fistulae.

[*5] Catheter was used for treatment and has been in place for 90 days or more prior to treatment. Patient does not have an fistula or graft in place. Catheter is only access. Port access devices are reported as catheters for this project.

reported as catheters for this project.

[*6] Patients listed as graft or catheter may have had fistulae in place for future use, but they actually received their treatment through a graft or catheter.

ADVANCED DIALYSIS CENTER - POTOMAC State: VA Network: 05 CCN: 492635

TABLE 8: Characteristics of New Dialysis Patients *1 , 2007-2010 (Form CMS-2728)

			This Fa	cility		Regional Averages *2, 2010			
	Measure Name	2007	2008	2009	2010	State	Network	U.S.	
Pat	ient Characteristics								
8a	Total number of patients with forms (n)	29	42	40	46	30.0	27.8	30.	
8b	Average age (years [0-95])*3	71.1	73.3	70.9	70.8	63.1	63.1	63.	
8c	Female (% of 8a)	44.8	42.9	40.0	52.2	45.1	44.8	43.	
8d	Race (% of 8a; sums to 100%)*4								
	African-American	55.2	35.7	32.5	41.3	45.3	47.3	27.	
	Asian/Pacific Islander	0.0	9.5	7.5	4.3	3.0	2.8	4.	
	Native American	0.0	2.4	0.0	0.0	0.1	0.2	1	
	White	44.8	52.4	60.0	54.3	51.4	49.6	66.	
	Other/Unknown/Missing	0.0	0.0	0.0	0.0	0.2	0.2	0.	
8e	Hispanic (% of 8a)	6.9	2.4	2.5	2.2	2.9	2.2	14.	
8f	Primary cause of ESRD (% of 8a; sums to 100%)								
	Diabetes	20.7	38.1	45.0	63.0	44.8	42.4	45.	
	Hypertension	65.5	54.8	50.0	28.3	31.5	34.2	29.	
	Primary Glomerulonephritis	0.0	0.0	0.0	0.0	6.1	5.9	7.	
	Other/Missing	13.8	7.1	5.0	8.7	17.5	17.6	18.	
8g	Medical coverage (% of 8a; sums to 100%)								
95	Employer group only	0.0	0.0	0.0	0.0	17.7	18.5	15.	
	Medicare only	62.1	21.4	17.5	13.0	24.6	23.1	23.	
	Medicaid only	3.4	0.0	12.5	2.2	7.6	10.7	11.	
	Medicare and Medicaid only	13.8	11.9	5.0	17.4	10.4	10.1	12.	
	Medicare and Other	6.9	42.9	47.5	45.7	22.7	21.9	21.	
	Other/Unknown	13.8	23.8	17.5	17.4	8.1	9.0	7.	
	None	0.0	0.0	0.0	4.3	8.9	6.6	7.	
8h	Body Mass Index *5								
OII	Male	22.5	23.9	27.4	27.2	28.1	27.9	27.	
	Female	27.8	33.3	26.3	28.8	30.1	29.6	28.	
0.		27.0	33.3	20.3	20.0	30.1	27.0	20.	
8i	Employment *6	4.5		500	20.0	20.0	20.2	2.4	
	Six months prior to ESRD treatment	16.7	66.7	50.0	20.0	39.0	38.3	34.	
	At first ESRD treatment	0.0	0.0	0.0	10.0	27.3	25.8	20.	
8j	Primary modality (% of 8a; sums to 100%)								
	Hemodialysis	100	100	100	100	93.9	93.1	93.	
	CAPD/CCPD	0.0	0.0	0.0	0.0	6.1	6.9	6.	
	Other/Unknown/Missing	0.0	0.0	0.0	0.0	0.0	0.0	0.	
8k	Number of incident hemodialysis patients (n)	29	42	40	46	28.4	26.0	27.	
81	Access used at first outpatient dialysis (% of 8k; sums to 100%)								
	Arteriovenous fistula	6.9	2.4	2.5	0.0	14.3	13.4	14.	
	Arteriovenous graft	0.0	4.8	2.5	0.0	3.5	3.3	2.	
	Catheter	93.1	92.9	95.0	100	82.1	83.1	81.	
	Other/Unknown/Missing	0.0	0.0	0.0	0.0	0.1	0.1	0.	
8m	Arteriovenous fistulae placed (% of 8k)	13.8	7.1	12.5	4.3	30.2	27.1	31.	
	erage Lab Values Prior to Dialysis *3	-2.0							
	Hemoglobin (g/dL [3-18])	9.9	10.3	0.6	0.5	0.6	0.6	0	
8n				9.6	9.5	9.6	9.6	9. 3.	
80	Serum Albumin (g/dL [0.8-6.0])	2.4	2.6	3.0	2.7	3.1	3.1		

(continued)

TABLE 8 (cont.): Characteristics of New Dialysis Patients*1, 2007-2010 (Form CMS-2728)

			This Fa	Regional Averages *2, 2010				
	Measure Name	2007	2008	2009	2010	State	Network	U.S.
Av	erage Lab Values Prior to Dialysis *3							
8p	Serum Creatinine (mg/dL [2-33])	4.8	4.0	5.2	5.1	6.3	6.3	6.2
8q	GFR (mL/min [0-60])	14.3	15.7	13.2	13.9	11.2	11.4	11.2
Ca	re Prior to ESRD Therapy							
8r	Received ESA prior to ESRD (% of 8a)	3.4	4.8	7.5	4.3	24.7	22.8	21.8
8s	Pre-ESRD nephrologist care (% of 8a; sums to 100%)							
	No	6.9	23.8	37.5	32.6	29.0	30.8	30.4
	Yes, < 6 months	3.4	2.4	2.5	2.2	14.3	15.3	13.2
	Yes, 6-12 months	0.0	14.3	20.0	28.3	19.0	18.7	19.1
	Yes, > 12 months	6.9	4.8	7.5	2.2	28.1	22.6	25.1
	Unknown/missing	82.8	54.8	32.5	34.8	9.6	12.6	12.2
8t	Informed of transplant options (% of 8a)	100	100	100	100	71.3	75.6	76.7
8u	Patients not informed of transplant options (n)	0	0	0	0	8.6	7.2	7.0
8v	Reason not informed (% of 8u; may not sum to 100%)							
	Medically unfit					17.3	18.0	31.4
	Unsuitable due to age					17.8	16.6	24.0
	Psychologically unfit					1.4	1.4	2.7
	Patient declined information					1.0	0.7	1.4
	Patient has not been assessed	•				60.3	62.4	44.5
Co	morbid Conditions							
8w	Pre-existing Comorbidity (% yes of 8a)							
	Congestive heart failure	27.6	38.1	30.0	32.6	30.7	30.1	32.2
	Atherosclerotic heart disease *7	3.4	2.4	2.5	0.0	15.5	15.0	20.8
	Other cardiac disorder *7	6.9	4.8	2.5	8.7	17.3	16.0	18.2
	CVD, CVA, TIA	20.7	4.8	0.0	10.9	9.8	8.7	9.4
	Peripheral vascular disease	6.9	0.0	0.0	6.5	9.1	10.0	13.6
	History of hypertension	82.8	81.0	97.5	89.1	85.2	84.8	86.2
	Diabetes *7	44.8	61.9	65.0	65.2	59.2	57.0	59.1
	Diabetes on insulin	27.6	42.9	57.5	60.9	39.5	37.0	38.2
	COPD	0.0	2.4	2.5	8.7	10.0	9.1	9.7
	Current smoker	3.4	2.4	0.0	4.3	7.3	6.1	6.3
	Cancer	0.0	0.0	2.5	8.7	7.0	6.2	7.7
	Alcohol dependence	0.0	0.0	0.0	0.0	1.9	1.6	1.8
	Drug dependence	0.0	0.0	0.0	0.0	0.9	1.1	1.3
	Inability to ambulate	0.0	2.4	0.0	2.2	6.6	5.7	7.4
	Inability to transfer	0.0	0.0	0.0	0.0	3.4	3.1	3.8
8x	Average number of comorbid conditions	2.2	2.4	2.6	3.0	3.0	2.9	3.2

n/a= not applicable

[*1] See *Guide, Section XI*.

[*2] Values are shown for the average facility.

[*3] For continuous variables, summaries include only responses in range indicated in brackets.

[*4] 'Asian' includes Indian sub-continent. 'Native American' includes Alaskan Native. 'White' includes Middle Eastern and Arabian.

[*5] The median BMI is computed for adult patients at least 20 years old.

[*6] Full-time, part-time, or student (% of 18-60 year olds).

^[47] Atherosclerotic heart disease' includes ischemic heart disease (coronary artery disease) and myocardial infarction. 'Other cardiac disorder' includes cardiac arrest, cardiac dysrhythmia, and pericarditis. 'Diabetes' includes patients with diabetes as the primary cause of ESRD.

TABLE 9: Summaries for All Dialysis Patients Treated as of December 31 of Each Year *1, 2007-2010

	This Facility Measure Name 2007 2008 2009 2010		Regiona	l Averages *2	, 2010			
	Measure Name	2007	2008	2009	2010	State	Network	U.S.
9a	Patients treated on 12/31 (n)	32	34	39	55	68.2	66.8	65.3
9b	Average age (years)	67.8	69.5	66.5	64.3	60.8	60.8	61.2
9c	Age (% of 9a; sums to 100%)							
	< 18	0.0	0.0	0.0	0.0	0.3	0.3	0.4
	18-64	40.6	32.4	46.2	45.5	57.7	57.6	56.2
	65+	59.4	67.6	53.8	54.5	42.0	42.1	43.4
9d	Female (% of 9a)	62.5	52.9	43.6	47.3	45.8	45.4	44.9
9e	Race (% of 9a; sums to 100%)							
	African American	40.6	55.9	64.1	61.8	59.0	61.3	37.3
	Asian/Pacific Islander	9.4	2.9	5.1	5.5	3.4	2.7	5.0
	Native American	0.0	2.9	2.6	1.8	0.2	0.2	1.4
	White	46.9	35.3	28.2	30.9	36.6	35.0	55.1
	Other/Unknown/Missing	3.1	2.9	0.0	0.0	0.8	0.8	1.
9f	Ethnicity (% of 9a; sums to 100%)							
	Hispanic	6.3	2.9	2.6	3.6	3.3	2.8	16.0
	Non-Hispanic	93.8	91.2	89.7	89.1	94.3	94.2	82.4
	Unknown	0.0	5.9	7.7	7.3	2.4	3.0	1.
9g	Cause of ESRD (% of 9a; sums to 100%)							
	Diabetes	46.9	50.0	51.3	56.4	40.5	39.1	44.1
	Hypertension	37.5	38.2	35.9	29.1	33.1	34.1	28.1
	Glomerulonephritis	0.0	0.0	0.0	0.0	11.8	11.3	11.5
	Other/Unknown	15.6	5.9	7.7	7.3	12.7	12.8	14.9
	Missing	0.0	5.9	5.1	7.3	1.9	2.7	1.3
9h	Average duration of ESRD (years)	2.9	2.9	3.9	4.5	4.6	4.6	4.5
9i	Years since start of ESRD (% of 9a; sums to 100%)							
	< 1	31.3	23.5	17.9	21.8	17.0	16.8	17.5
	1-2	21.9	20.6	25.6	12.7	18.4	18.3	18.4
	2-3	15.6	17.6	7.7	9.1	13.6	13.8	14.2
	3-6	18.8	26.5	23.1	29.1	26.7	26.2	26.3
	6+	12.5	11.8	25.6	27.3	24.3	24.9	23.6
9j	Nursing home patients (% of 9a) *3	90.6	94.1	82.1	74.5	12.4	14.0	12.7
9k	Modality (% of 9a; sums to 100%)							
	In-center hemodialysis	93.8	100	100	96.4	89.7	90.0	89.9
	Home hemodialysis	0.0	0.0	0.0	0.0	1.7	1.3	1.5
	Continuous ambulatory peritoneal dialysis	0.0	0.0	0.0	0.0	2.6	2.5	2.3
	Continuous cycling peritoneal dialysis	0.0	0.0	0.0	0.0	5.1	5.1	5.0
	Other modality *4	6.3	0.0	0.0	3.6	0.9	1.0	0.7

n/a = not applicable
[*1] See *Guide, Section XII.*[*2] Values are shown for the average facility.
[*3] Includes patients who were also treated by a nursing facility at any time during the year. The source of nursing facility history of patients is the Nursing Home Minimum Dataset.
[*4] Other modality includes other dialysis, uncertain modality, and patients not on dialysis but still temporarily assigned to the facility (discontinued dialysis, recovered renal function, and lost to follow up).

TABLE 10: Comorbidities Reported on Medicare Claims for Medicare Dialysis Patients Treated as of December 31 of Each Year *1, 2007-2009

		Th	is Facility	7	Regiona	l Averages *2	, 2009
	Measure Name	2007	2008	2009	State	Network	U.S.
10a	Medicare dialysis patients on 12/31 (n)	26	31	30	52.8	52.5	50.8
10b	Comorbidity (% yes of 10a)						
	AIDS/HIV positive	0.0	3.2	0.0	1.7	3.1	1.8
	Alcohol dependence	3.8	9.7	6.7	2.0	2.1	2.0
	Anemia	11.5	19.4	20.0	6.3	6.5	6.6
	Cancer	15.4	12.9	13.3	11.8	11.9	10.8
	Cardiac arrest	3.8	6.5	13.3	1.6	1.5	1.5
	Cardiac dysrhythmia	50.0	67.7	53.3	33.2	38.4	34.5
	Cerebrovascular disease	76.9	67.7	63.3	26.3	28.2	26.6
	Chronic obstructive pulmonary disease	50.0	38.7	43.3	27.2	28.1	29.1
	Congestive heart failure	84.6	87.1	73.3	51.6	52.3	50.8
	Diabetes	80.8	90.3	90.0	61.0	61.1	62.2
	Drug dependence	0.0	0.0	0.0	1.9	2.1	1.9
	Gastrointestinal tract bleeding	15.4	6.5	10.0	2.9	2.9	2.7
	Hepatitis B	0.0	0.0	0.0	0.8	1.2	1.1
	Hepatitis other	7.7	6.5	6.7	4.1	6.3	4.8
	Hyperparathyroidism	3.8	9.7	20.0	11.3	8.7	8.2
	Infection: dialysis access-related	73.1	58.1	46.7	18.9	19.0	18.9
	Infection: not related to dialysis access	96.2	87.1	83.3	47.3	51.8	49.8
	Ischemic heart disease	76.9	77.4	63.3	47.0	48.5	49.9
	Myocardial infarction	15.4	29.0	26.7	8.5	9.0	8.4
	Peripheral vascular disease *3	57.7	77.4	63.3	40.1	44.1	44.3
	Pneumonia	34.6	19.4	26.7	5.4	5.6	5.6
10c	Average number of comorbid conditions	7.6	7.7	7.2	4.1	4.3	4.2

n/a = not applicable
[*1] Based on patients with Medicare as primary insurer on 12/31 each year. See *Guide, Section XIII*.
[*2] Values are shown for the average facility.
[*3] Peripheral vascular disease includes both venous and arterial diseases.

TABLE 11: How Patients Were Assigned to This Facility and End of Year Patient Status *1, 2007-2010

		This Facility			Regional Averages *2, 2010			
	Measure Name	2007	2008	2009	2010	State	Network	U.S.
11a	Number of patients placed in facility *1 (n)	73	93	100	133	93.4	94.1	90.9
11b	Initial patient placement for the year in this facility							
	(% of 11a; sums to 100%)							
	Continuing at facility on 01/01	24.7	34.4	34.0	29.3	70.3	69.0	69.0
	Incident (new to ESRD)	27.4	25.8	22.0	25.6	19.3	18.8	19.3
	Transferred into facility	47.9	39.8	44.0	45.1	10.5	12.2	11.7
11c	Patient status at end of year (% of 11a; sums to 100%)							
	Alive in this facility on 12/31	43.8	36.6	39.0	41.4	73.0	70.9	71.9
	Alive in another facility on 12/31	21.9	18.3	15.0	24.1	6.8	8.4	8.4
	Received a transplant	0.0	0.0	0.0	0.0	2.9	2.9	2.7
	Died; death attributed to this facility	24.7	32.3	35.0	26.3	13.7	13.4	13.5
	Died; death attributed to another facility	6.8	8.6	6.0	3.8	1.3	1.8	1.4
	Other*3	2.7	4.3	5.0	4.5	2.3	2.6	2.2

^[*1] Patient assignment for Tables 1,2,3,9, and 10 only. See *Guide, Section XIV*.
[*2] Values are shown for the average facility.
[*3] Includes patients who recovered renal function, discontinued dialysis, or were lost to follow-up. Also includes dialysis unrelated deaths. Dialysis unrelated deaths are not attributed to any facility for the purposes of the mortality calculations in this report.

TABLE 12: Patient and Staff Counts from the Annual Facility Survey (Form CMS-2744)*1, 2007-2010

		This Facility			Regional Averages *2, 2010			
	Measure Name	2007	2008	2009	2010	State	Network	U.S.
Pat	ients Treated During the Year							
12a	Patients treated during year (n)	137	177	196	220	107.2	108.5	103.0
12b	Incident patients (% of 12a)	26.3	23.2	24.0	25.5	19.5	18.6	19.
12c	Transferred into facility (% of 12a)	48.9	52.5	50.5	47.7	12.7	15.5	13.
12d	Transferred out of facility (% of 12a)	40.1	42.4	41.3	39.1	13.0	16.0	14.
Pat	ients Treated on 12/31							
12e	Patients treated on 12/31 (n)	50	48	59	71	73.4	71.8	70.
12f	Patient modality on 12/31 (n; sums to 12e)							
	In-center HD	50	48	59	71	66.6	65.5	64.
	In-center CAPD	0	0	0	0	0.0	0.0	0.
	In-center CCPD	0	0	0	0	0.0	0.0	0.
	In-center Other	0	0	0	0	0.0	0.0	0.
	Home HD	0	0	0	0	1.2	0.9	1.
	Home CAPD	0	0	0	0	2.1	1.9	1.
	Home CCPD	0	0	0	0	3.5	3.5	3.
	Home Other	0	0	0	0	0.0	0.0	0.
2g	Medicare eligibility status on 12/31 (% of 12e; sums to 100% *3)							
	Medicare	74.0	89.6	91.5	91.5	88.9	86.6	86.
	Medicare application pending	12.0	2.1	1.7	2.8	6.8	8.1	9.
	Non-Medicare	14.0	8.3	6.8	5.6	4.3	5.2	4.
Fac	cility Staffing on 12/31 *4							
l2h	Total full and part time staff positions (n)	12	12	12	16	14.4	14.6	14.
12i	Staff positions by type (n; sums to 12h)							
	Full time nurse *5	2	2	2	3	5.0	4.6	4.
	Full time patient care technician	6	5	6	6	5.4	6.1	5.
	Full time renal dietician	1	0	0	0	0.5	0.5	0
	Full time social worker	1	0	0	0	0.5	0.5	0
	Part time nurse *5	1	1	0	3	0.9	0.9	1.
	Part time patient care technician	1	2	2	2	0.9	0.9	0.
	Part time renal dietician	0	1	1	1	0.6	0.6	0.
	Part time social worker	0	1	1	1	0.6	0.6	0.

^[*1] See *Guide, Section XV*.
[*2] Values are shown for the average facility.
[*3] Values may not sum to exactly 100% because of unknown Medicare status.
[*4] Data as of July 31, 2011. A *full time position* is defined as a position with at least 32 hours of employment a week and a *part time position* is defined as a position with less than 32 hours of employment a week (includes positions that were opened but not filled on this date).
[*5] Nursing staff includes registered nurse, licensed practical nurse, vocational nurse, or advanced practice nurse degree.

TABLE 13: Survey and Certification Activity*1, October 2008-June 2011

		This Facility	Regional Averages *2, October 2008-June 2011			
	Measure Name	Survey	State	Network	U.S.	
13a	Date of last survey	02/05/2009	n/a	n/a	n/a	
13b	Type of last survey	RECERTIFICATION	n/a	n/a	n/a	
13c	Compliance condition after last survey	Meets requirements	n/a	n/a	n/a	
13d	Number of deficiencies cited at last survey *3					
	Condition for coverage (CfC) deficiencies	0	0.1	0.1	0.4	
	Standard deficiencies	2	2.6	7.3	7.3	
13e	CfC deficiencies cited at last survey *3					
	V100 Compliance with Fed., State, and Local	No, not cited	0.0	0.0	0.1	
	V110 Infection Control	No, not cited	1.7	3.1	5.4	
	V175 Water and Dialysate Quality	No, not cited	1.7	1.1	4.3	
	V300 Reuse of Hemodialysis and Bloodlines	No, not cited	0.0	0.0	0.4	
	V400 Physical Environment	No, not cited	0.9	1.1	2.5	
	V450 Patient Rights	No, not cited	0.0	0.0	0.4	
	V500 Patient Assessment.	No, not cited	0.0	1.5	4.0	
	V540 Patient Plan of Care	No, not cited	1.7	2.3	4.7	
	V580 Care at Home	No, not cited	0.0	0.0	0.8	
	V625 Quality Assessment & Performance improvement	No, not cited	0.9	1.9	4.2	
	V660 Special Purpose Renal Dialysis Facilities	No, not cited	0.0	0.0	0.0	
	V675 Laboratory Services	No, not cited	0.0	0.0	0.1	
	V680 Personnel Qualifications	No, not cited	0.0	0.0	0.9	
	V710 Responsibilities of the Medical Director	No, not cited	0.9	0.4	4.7	
	V725 Medical Records	No, not cited	0.0	0.0	0.5	
	V759 Governance	No, not cited	1.7	2.3	4.6	

TABLE 14: Facility Information *1 , 2011

Field	This Facility
Ownership:	For Profit
Organization:	Not applicable
Initial Medicare certification date:	06/21/2006
Number of stations *2:	9
Services provided *2:	Unavailable
CMS Certification Numbers (CCN) included in this report:	492635
National Provider Identifier (NPI) *3:	1811095722

n/a = not applicable
[*1] See *Guide, Section XVI.*[*2] Average values are shown for the latest survey at each facility during the period October 2008 through June 2011. Some surveys from April through June of 2011 may not be included.
[*3] Values are shown for the latest survey at each facility during the period October 2008 through June 2011.

^[*1] Information based on SIMS data as of March 31, 2011. See *Guide, Section XVII*.
[*2] Information based on data available on DFC as of Aug 2, 2011.
[*3] Information based on CROWNWEB data as of April 2011. If missing, data was not available.